

The National STD Curriculum

Official Launch Webinar

David Spach, M.D.

Professor of Medicine, Division of Infectious Diseases
University of Washington

Introductions By:

Jami Frazee, PhD, NNPTC Project Officer
Gail Bolan, MD, DSTDP Division Director

January 25, 2018



PROVIDERS

ARE

PIVOTAL

Gail Bolan, MD

Director, Division of STD Prevention
Centers for Disease Control and Prevention

Visit: <https://www.cdc.gov/std/default.htm>

2018 STD PREVENTION
CONFERENCE

Call for Abstracts OPEN through Feb. 16, 2018

“intersecting epidemics, integrated solutions”





Centers for Disease Control and Prevention

Resources

- **STD Treatment Guidelines**

- Visit: www.cdc.gov/std/tg2015/default.htm

- **FREE STD Treatment Guidelines App (Apple and Android)**



National Network of STD Clinical Prevention Training Centers (NNPTC)



Resources

GOT A TOUGH STD QUESTION?
Get FREE expert STD clinical consultation at your fingertips

Ask your question

National STD experts review

Response within 1-5 business days, depending on urgency

Log on to www.STDCCN.org for medical professionals nationwide

- Visit: www.nnptc.org
- Clinical Training and Consultation Network
 - Visit: www.STDCCN.org.
- Resources and tools for STD treatment
- STD Clinical Toolbox App



National **STD**
Curriculum

National STD Curriculum

David H. Spach, MD
Professor of Medicine
Division of Infectious Diseases
University of Washington

January 25, 2018


Disclosures: NONE



National STD Curriculum

 Contributors

Funded by a grant from the
Centers for Disease Control and Prevention



Objectives

- Provide an overview of the *National STD Curriculum* web site
- How to optimize use of the curriculum
- Explain administration functions (register, track progress, CE)
- Summarize “1 Year” site data
- Describe future activities on site



STD Modules Self-Study >



STD Modules Quick Reference >



Question Bank



Clinical Consultation




Master Bibliography



National STD Curriculum

 Contributors

Funded by a grant from the Centers for Disease Control and Prevention



This project is funded by a grant from the Centers for Disease Control and Prevention

National STD Curriculum: *Acknowledgments*

- **Editors**

- David Spach
- Jeanne MARRAZZO
- Christine Johnston
- Kimberly Workowski

- **Technology Team**

- Andrew Karpenko
- Kenton Unruh

- **Project Management**

- Bruce Maeder
- Julia Freimund
- Mary Annese
- Karin Bauer

- **CDC Division of STD Prevention**

- Diane Ballard
- Roxanne Barrow
- Gail Bolan
- Jami Frazee
- Jennifer Fuld
- Suzanne Haecker
- Sheila McKenzie
- Rachel Powell
- Salina Smith

- **Continuing Education Partners**

- CME at UAB
- CNE at UW School of Nursing

National STD Curriculum: *Acknowledgments*

- **Additional Authors**

- Elfriede A. Agyemang
- Lindley A. Barbee
- William M. Geisler
- Seth R. Glassman
- Andrew W. Hahn
- Rebecca G. Kinney
- Laura A. Quilter

- **Additional Reviewers**

- Khalil G. Ghanem
- Margaret R. Hammerschlag
- Katherine Hsu
- Hillary Liss
- Ina Park
- Nicholas Van Wagoner

Note: most of the authors and reviewers are part of the
National Network of STD Clinical Prevention Training Centers

NATIONAL STD CURRICULUM

Site Overview



STD Modules
Self-Study >



STD Modules
Quick Reference >



Question
Bank



Clinical
Consultation




Master
Bibliography



National STD Curriculum

Contributors

Funded by a grant from the
Centers for Disease Control and Prevention



To access the *National STD Curriculum* go to:
www.std.uw.edu

National STD Curriculum: Features

- Interactive self-paced on-line learning and assessment tool
- Free CNE and CME credits
- Expert, up-to-date content consistent with CDC guidelines
- Free module-by-module certificate program
- Group functionality


Intended audience: Medical Professionals




National STD Curriculum

Contributors

Funded by a grant from the Centers for Disease Control and Prevention



STD Modules



Chlamydia

[Chlamydia Self-Study](#) **CNE/CME**

Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about Chlamydia

[Question Bank](#) **CNE/CME**

Interactive board-review style questions with CE credit



Gonorrhea

[Gonorrhea Self-Study](#) **CNE/CME**


Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about Gonorrhea

[Question Bank](#) **CNE/CME**

Interactive board-review style questions with CE credit



HSV

Herpes Simplex Virus (HSV)

[HSV Self-Study](#) **CNE/CME**


Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about HSV

[Question Bank](#) **CNE/CME**

Interactive board-review style questions with CE credit



HPV

Human Papillomavirus (HPV)

[HPV Self-Study](#) **CNE/CME**

Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about HPV



PID

Pelvic Inflammatory Disease (PID)

[PID Self-Study](#) **CNE/CME**

Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about PID



Syphilis

[Syphilis Self-Study](#) **CNE/CME**

Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about Syphilis



Vaginitis

[Vaginitis Self-Study](#) **CNE/CME**

Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about Vaginitis

STD Modules



STD Modules Self-Study >



STD Modules Quick Reference >



Question Bank



Clinical Consultation




Master Bibliography




National STD Curriculum

Contributors

Funded by a grant from the Centers for Disease Control and Prevention



STD Modules



Chlamydia

[Chlamydia Self-Study](#) **CNE/CME**

Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about Chlamydia

[Question Bank](#) **CNE/CME**

Interactive board-review style questions with CE credit



Gonorrhea

[Gonorrhea Self-Study](#) **CNE/CME**


Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about Gonorrhea

[Question Bank](#) **CNE/CME**

Interactive board-review style questions with CE credit



HSV

Herpes Simplex Virus (HSV)

[HSV Self-Study](#) **CNE/CME**


Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about HSV

[Question Bank](#) **CNE/CME**

Interactive board-review style questions with CE credit



HPV

Human Papillomavirus (HPV)

[HPV Self-Study](#) **CNE/CME**

Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about HPV



PID

Pelvic Inflammatory Disease (PID)

[PID Self-Study](#) **CNE/CME**

Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about PID



Syphilis

[Syphilis Self-Study](#) **CNE/CME**

Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about Syphilis



Vaginitis

[Vaginitis Self-Study](#) **CNE/CME**


Track progress and receive CE credit

[Quick Reference](#) >


Rapidly access info about Vaginitis

STD Modules
Self-StudySTD Modules
Quick ReferenceQuestion
BankClinical
ConsultationMaster
Bibliography

National STD Curriculum

[Contributors](#)Funded by a grant from the
Centers for Disease Control and Prevention

STD Modules



Chlamydia

[Chlamydia Self-Study](#) **CNE/CME**

Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about Chlamydia

[Question Bank](#) **CNE/CME**Interactive board-review style
questions with CE credit


Gonorrhea

[Gonorrhea Self-Study](#) **CNE/CME**

Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about Gonorrhea

[Question Bank](#) **CNE/CME**Interactive board-review style
questions with CE credit


HSV

Herpes Simplex Virus
(HSV)[HSV Self-Study](#) **CNE/CME**

Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about HSV

[Question Bank](#) **CNE/CME**Interactive board-review style
questions with CE credit

HPV

Human Papillomavirus
(HPV)[HPV Self-Study](#) **CNE/CME**

Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about HPV



PID

Pelvic Inflammatory
Disease (PID)[PID Self-Study](#) **CNE/CME**

Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about PID



Syphilis

[Syphilis Self-Study](#) **CNE/CME**

Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about Syphilis



Vaginitis

[Vaginitis Self-Study](#) **CNE/CME**

Track progress and receive CE credit

[Quick Reference](#) >

Rapidly access info about Vaginitis

STD Modules: Dual Functionality

Self-Study

- Sequential (Step-by-Step)
- Flexible modular options
- Certificate program
- Ideal for training programs

Quick Reference

- Highly organized interface
- Quick search
- On demand topics
- Ideal for staying updated



STD Modules Self-Study >


STD Modules Quick Reference >

Question Bank


Clinical Consultation

Master Bibliography


Self-Study STD Modules




Chlamydia




Gonorrhea




HSV




HPV



PID



Syphilis



Vaginitis



Self-Study STD Modules for Clinicians

You are just a few steps away from free CE credits!

1 ★

Sign in or Register

A free account is required.

2

Study the Materials

An entire module, or just a few topics at a time.

3

Take the CE Quiz

5 questions covering the topics in each module.

4

Claim CE Credit

Free CNE and Free CME available!

New Users

Create a free account to get started.

Required for CE

Register >

Returning Users

Email Address

Password

[Forgot password?](#)

Sign in >



Self-Study STD Modules



Chlamydia

Gonorrhea

HSV

HPV

PID

Syphilis

Vaginitis

Certificate of Completion

Module 6. Syphilis



Syphilis Overview

Back

Next

About this Module

The **Syphilis STD Module** is a web-based training course designed to guide clinicians in the diagnosis, treatment, and prevention of syphilis. This module is based on original content developed by the National Network of STD Clinical Prevention Training and incorporates recommendations from the 2015 CDC STD Treatment Guidelines. The original online Syphilis STD Module was developed by the Division of STD Prevention, Centers for Disease Control and Prevention.

Last Updated: December 21st, 2017

CNE/CME Continuing Education

This module qualifies for:

- 1 CME AMA PRA Category 1 Credits™, or
- 1 CNE contact hour

[View CE Notices](#)

CNE and CME Origination: February 1st, 2017

CNE and CME Expiration: January 31st, 2020

Steps to Acquire CE for this Activity:

1 ★

CE Quiz
Score 80%+

2

Give Feedback
Complete survey

3

Print Certificate
Obtain proof of CE

Target Audience

The **Syphilis** Self-Study STD Module is intended for the following clinicians who desire a basic introduction to STD diagnosis and management:

- Physicians
- Physician assistants
- Nurse practitioners
- Nurses
- Nurse-midwives

Lesson Plan

1

Topic 1
Background

2

Topic 2
Epidemiology in the United States

3

Topic 3
Clinical Manifestations

4

Topic 4
Treatment

5

Topic 5
Microbiology and Pathogenesis

6

Topic 6
Laboratory Diagnosis

7

Topic 7
Screening for Infection

Self-Study STD Modules



Chlamydia



Gonorrhea



HSV



HPV



PID



Syphilis

Vaginitis



Certificate of Completion

Topic 1. Background

📄 Topic Objective

🔍 Back

Next



BACKGROUND

Syphilis is a systemic infection caused by *Treponema pallidum*, a spirochete bacterium that is transmitted primarily through sexual activity. In the absence of treatment, patients who acquire *T. pallidum* remain chronically infected and syphilis generally progresses in stages, characterized by episodes of active clinical manifestations interrupted by periods of latent infection. Chronic disease can result in significant morbidity, potentially affecting nearly every organ system, and rarely, can result in death. In addition, untreated syphilis in pregnant women can lead to fetal demise and devastating congenital infection for neonates born to an infected mother.

🔍 Back

Next



Topic 2. Epidemiology in the United States

Topic Objective

Back

Next

EPIDEMIOLOGY IN THE UNITED STATES

INCIDENCE TRENDS IN THE UNITED STATES

In the United States, the reporting of syphilis incidence is usually represented in terms of primary and secondary syphilis.^[1] Reported cases of primary and secondary syphilis (P & S syphilis) represent the incidence of syphilis more accurately than reported cases of latent infection, particularly late latent syphilis, which signifies infection acquired more than a year before syphilis is diagnosed. In the past 76 years, in the United States, the syphilis epidemic has undergone major fluctuations (Figure 1).^[1] During the 1920s and 1930s, syphilis rates were very high, but declined rapidly in the late 1940s with the introduction of penicillin.^[1,2] The number of reported cases rose between 1986 and 1990, but by the late 1990s, syphilis rates in the United States had declined to a point where public health authorities declared syphilis elimination a feasible goal and in 1999 the CDC developed a national plan to eliminate syphilis in the United States.^[3] In 2000, the reported rate of syphilis in the United States reached an all-time low, but unfortunately, cases of syphilis have increased since 2001 (Figure 2), primarily due to an increase among men who have sex with men (MSM), including those coinfected with HIV.^[1] In 2016, a total of 27,814 cases of primary and secondary syphilis were reported, which corresponds to a rate of 8.7 cases per 100,000 population; the total number cases of primary and secondary syphilis in 2016 represent a 16.5% increase from 2015, a 102% increase from 2010, and the highest reported number of cases since 1994.^[1]

EPIDEMIOLOGY BASED ON SPECIFIC DEMOGRAPHICS

The following summarizes syphilis epidemiology in the United States based on demographics per the 2016 CDC STD Surveillance report.^[1]

Geographic Distribution

In the United States, from 2012 to 2016, the rates of primary and secondary syphilis




Figure 1. Syphilis Rates by Stage of Infection, United States, 1941-2016




Figure 2. Syphilis Cases, All Stages of Infection, United States, 2000-2016

Check on Learning (optional)

NSTDC | STD Modules Self-Study | STD Modules Quick Reference | Question Bank | Clinical Consultation | Master Bibliography

Based on National Health and Nutrition Examination Surveys (NHANES), chlamydia prevalence in the U.S. is estimated to be 1.5%.^[4] Chlamydia prevalence is highest among adolescents and young adults, as well as among racial and ethnic minorities. Test positivity is often used as a proxy of chlamydia prevalence in a population. During 2007 to 2012, chlamydia test positivity among males and females aged 14 to 39 years was 1.7%.^[1] Among sexually active females aged 14 to 24 years (the population targeted for routine screening), chlamydia prevalence was 4.7%; black females had, by far, the highest prevalence (Figure 7).^[1]

RISK FACTORS

Risk factors associated with acquisition of chlamydial infection include new or multiple sex partners, a history of STIs, presence of another STI, and lack of barrier contraception.^[5] The presence of columnar epithelial cells on the ectocervix, referred to as ectopy, is a condition that may increase susceptibility to chlamydial infection; oral contraceptive use contributes to ectopy.^[6] Adolescents and young adults are at increased risk for chlamydial infection for a combination of biological, behavioral, and cultural reasons, including difficulty accessing preventive health care services for STIs.

Figure 2. Chlamydia—Rate, by Sex, 2000-2016



Figure 3. Chlamydia—Rates of Reported Cases by Age Group and Sex, U.S. 2016




Figure 2. Chlamydia—Rate, by Sex, 2000-2016




Figure 3. Chlamydia—Rates of Reported Cases by Age Group and Sex, U.S. 2016



Chlamydia—Rates of Reported Cases by Race/Ethnicity, U.S. 2016



Check On Learning Question

Based on United States STD surveillance data, which age group has the highest rates of infection with *Chlamydia trachomatis*?

- Males aged 15-24 years
- Males aged 30-39 years
- Females aged 15-24 years
- Females aged 30-39 years

Check On Learning Question







Based on United States STD surveillance data, which age group has the highest rates of infection with *Chlamydia trachomatis*?

You chose this option correctly:

- Females aged 15-24 years

Females 15-24 years of age are the age group with the highest rates of *C. trachomatis* infection. The high rates in this age group, in combination with the significant potential for long-term complications in women from unidentified *C. trachomatis* infection, underlies the USPSTF and CDC recommendation to perform routine screening in all sexually active women younger than 25 years of age.

Sample Illustrations on Web Site



Primary Genital Infection

Primary infection is defined as the first infection with either HSV-1 or HSV-2 with absence of antibody to either HSV type. Primary genital infection is often symptomatic, but patients may have unrecognized or subclinical infection. With symptomatic infection, clinical manifestations of primary infection typically resolve within 3 weeks in the absence of antiviral therapy. Serum antibodies appear within 12 weeks of the primary infection in most persons.^[54] The following symptoms may occur with primary HSV-1 or HSV-2 genital infection:

- Severe multiple bilateral genital ulcers, pain, itching, dysuria, vaginal or urethral discharge, and tender inguinal adenopathy (Figure 7 [Graphic Image](#)).
- Without antiviral therapy, lesions last 2 to 3 weeks, with evolution of the lesions from vesicle pustule to wet ulcers to dry crusts (Figure 8).^[55]
- The median duration of viral shedding is about 10 to 12 days, and correlates with the time from the onset of vesicles to crusting of lesions.^[56]
- Systemic symptoms (fever, myalgias, headaches, aseptic meningitis or symptoms of autonomic nervous system dysfunction such as urinary retention) peak within 3 to 5 days of onset of lesions and gradually recede over the next 3 to 4 days.^[57]
- In women, HSV shedding from the cervix occurs in 80 to 90% of primary HSV-1 and HSV-2 infections.^[52] Cervicitis may involve the ectocervix or endocervix, with or without clinical symptoms. In most cases, the cervix appears abnormal to inspection with ulcerative lesions, erythema, or friability.
- Herpes proctitis typically manifests with fever, pain, discharge, tenesmus, and constipation; some patients will have severe anal ulcerations visible on anoscopy; some patients develop symptoms of autonomic dysfunction, including difficulty urinating.^[58] Rarely, herpes proctitis may present as a pseudotumor that mimics epidermoid carcinoma.^[59]
- Infection of the urethritis and/or meatus may cause a clear mucoid discharge.^[60]

Nonprimary Infection

The term nonprimary HSV infection most often refers to infection with HSV-1 or HSV-2 in an individual with pre-existing antibodies to the other virus. For example, a person may acquire oral HSV-1 infection as a child and later acquire genital HSV-2 as an adult. Manifestations of nonprimary infection tend to be milder than those of primary infection, presumably due to cross-immunity protection from prior infection with the other HSV type.^[61,62] Less often, in a different scenario, nonprimary infection can




Figure 7. Primary Genital HSV Infection




Figure 8. Clinical and Virologic Course of Genital HSV in Primary Infection



data demonstrate that more widespread resistance has emerged with some antimicrobials and might develop in the near future with others, thus highlighting the need for ongoing surveillance. The GISP tracks primary antimicrobial drugs used to treat gonorrhea in the United States and *N. gonorrhoeae* susceptibility to 7 antimicrobials: ceftriaxone, cefixime, azithromycin, spectinomycin, ciprofloxacin, penicillin, and tetracycline (Figure 8).^[1,3] These specific drugs are tested because they either **Rocephin** recently or were previously used for gonorrhea treatment.

- Ceftriaxone:** During 2006–2016, the percentage of isolates with reduced ceftriaxone susceptibility (defined as MIC ≥ 0.125 $\mu\text{g}/\text{mL}$) fluctuated between 0.05% and 0.4%.^[1] In 2016, the percentage of isolates with reduced susceptibility to ceftriaxone (MICs ≥ 0.125 $\mu\text{g}/\text{mL}$) was approximately 0.3%. In the GISP program, five isolates have been reported with a ceftriaxone MIC of 0.5 $\mu\text{g}/\text{mL}$.^[1]
- Cefixime:** The proportion of *N. gonorrhoeae* isolates in the United States with elevated cefixime minimum inhibitory concentrations (≥ 0.25 $\mu\text{g}/\text{mL}$) increased from 2009 to 2011, fell in 2012 and 2013, increased in 2014, and decreased in 2015 and 2016.^[1] In 2016, the percentage of elevated cefixime MICs (≥ 0.25 $\mu\text{g}/\text{mL}$) was approximately 0.3%, which is a significant decline from the peak of 1.4% in 2010 and 2011.^[1]
- Azithromycin:** Gonococcal azithromycin resistance has been tracked since 1992. From 2012–2016, the percentage of isolates with reduced azithromycin susceptibility (MICs ≥ 2 $\mu\text{g}/\text{mL}$) ranged from 0.02% to 3.6%; between 2014 and 2016, the percentage increased from 2.4% to 3.6%.^[1]

Ciprofloxacin: Fluoroquinolone-resistant *N. gonorrhoeae* is widely disseminated




Figure 5. Gonorrhea — Rates of Reported Cases by Sex, United States, 2012–2016




Figure 6. Gonorrhea — Rates of Reported Cases by Race/Ethnicity, 2016



STD Modules Self-Study >



STD Modules Quick Reference



Question Bank



Clinical Consultation



Master Bibliography

For quick access viewing of content in Self-Study

Chlamydia >

Gonorrhea >

Herpes Simplex Virus (HSV) >

Human Papillomavirus (HPV) >

Pelvic Inflammatory Disease (PID) >

Syphilis >

Vaginitis >

Chlamydia Quick Reference

TOPICS

1. Epidemiology in the United States
2. Microbiology and Pathogenesis
3. Clinical Manifestations
4. Laboratory Diagnosis
5. Screening for Chlamydial Infection
6. Treatment
7. Patient Counseling and Education
8. Summary Points


CNE/CME Chlamydia Self-Study Module

National

Contributors


Funded by a grant from the Centers for Disease Control and Prevention

STD Modules



Chlamydia

Track progress



Gonorrhea

Gonorrhea Self-Study CNE/CME
Track progress and receive CE credit

Chlamydia

CME/CNE Self-Study Module PDF Share

Last Updated: December 20th, 2017
Authors: Andrew W. Hahn, MD, William M. Geisler, MD, MPH
Reviewers: Kimberly A. Workowski, MD, Margaret R. Hammerschlag, MD

EPIDEMIOLOGY IN THE UNITED STATES

INCIDENCE

Chlamydia is the most common reportable bacterial sexually transmitted infection (STI) in the United States, with 1,598,354 cases reported in 2016.^[1] Since many persons with chlamydial infection may have minimal or no symptoms, the actual number of annual infections is significantly higher than the reported cases.^[2] The number of reported chlamydia cases have significantly increased since the early years of reporting that began in the 1980's (Figure 1),^[1] which may reflect an increase in the number of true infections, enhanced screening with more sensitive diagnostic tests, or a combination of both. Chlamydial rates of reported cases have consistently been higher in women than in men (Figure 2), with the highest rates (reported cases per 100,000 population) among females 15 to 24 years of age (Figure 3).^[3] In the United States, racial and ethnic minorities are disproportionately affected by chlamydia, particularly blacks (Figure 4).^[1] Factors contributing to these inequities may include differential access to quality health care, social and economic conditions, higher prevalence of disease in sexual networks, and differences in immunogenetic determinants that influence the immune response to chlamydia. The South has consistently had the highest rate of reported chlamydia cases, although the difference

Table of Contents
Chlamydia
Epidemiology in the United States
Microbiology and Pathogenesis
Clinical Manifestations
Laboratory Diagnosis
Screening for Chlamydial Infection
Treatment
Patient Counseling and Education
Summary Points
<input checked="" type="checkbox"/> Check-On-Learning
Citations
Additional References
Figures
Tables




Figure 1. Chlamydia Cases in U.S., 1980-2016





LABORATORY DIAGNOSIS

The selection of a laboratory test to detect the presence of *C. trachomatis* is a critical component of disease management and prevention.^[36] The testing technology has shifted from culture-based methods to molecular-based techniques and this represents a substantial improvement in test sensitivity and ease of specimen collection.

NUCLEIC ACID AMPLIFICATION TESTS (NAATS)

Nucleic acid amplification tests (NAATs) amplify nucleic acid sequences (either DNA or RNA) that are specific for the organism being detected. Similar to other nonculture tests, NAATs can detect live or non-viable organisms. *C. trachomatis* NAATs are FDA-cleared for use on urine specimens from men and women, urethral swabs in men, and endocervical swabs in women; some tests are cleared for vaginal swabs.^[2] The use of *C. trachomatis* NAAT for pharyngeal and rectal specimens is not FDA approved; however, laboratories can perform certain validation procedures, such as Clinical Laboratory Improvement Amendment (CLIA)-defined performance specifications, to enable them to test specimens for clinical purposes. In men, NAATs are the most sensitive and recommended test for detecting *C. trachomatis* from a urethral swab or first-catch urine specimen.^[2] For chlamydia screening in women, vaginal swabs are preferred over urine samples and several studies have shown that self-collected vaginal swabs are preferred by women and perform equal to or better than clinician-collected vaginal swabs.^[40,41,42,43,44] In addition, in men and women, self-collected rectal swabs for NAAT have also performed well.^[45] There is currently insufficient evidence to support the use of self-collected oropharyngeal or penile meatal swabs for the diagnosis of chlamydia.^[2] Multiple NAATs are commercially available for the detection of *C. trachomatis*.

NON-AMPLIFICATION MOLECULAR TESTS

Molecular tests that do not use nucleic acid amplification encompass a variety of antigen detection and nucleic acid hybridization methods. These include enzyme-immunoassays (EIA), direct fluorescent antibody tests (DFA), and nucleic acid hybridization tests, a distinct non-NAAT methodology that detects *C. trachomatis*-specific DNA or RNA sequences in rRNA, genomic DNA, or plasmid DNA. All have significantly

Table of Contents

Chlamydia

[Epidemiology in the United States](#)[Microbiology and Pathogenesis](#)[Clinical Manifestations](#)[Laboratory Diagnosis](#)[Nucleic Acid Amplification Tests \(NAATs\)](#)[Non-amplification Molecular Tests](#)[Culture](#)[Serology](#)[Diagnostic Evaluation in Suspected Sexual Abuse in Children](#)[Reporting Requirements](#)[Screening for Chlamydia Infection](#)[Treatment](#)[Patient Counseling and Education](#)[Summary Points](#)[Check-On-Learning](#)

Site Overview

Question Bank

STD Modules
Self-StudySTD Modules
Quick ReferenceQuestion
BankClinical
ConsultationMaster
Bibliography

STD Question Bank

Pathogen-Based
Diseases

Bacterial Vaginosis

Candidiasis – Vulvovaginal

Chancroid

Chlamydia

Gonorrhea

Granuloma Inguinale

Herpes Simplex Virus – Genital

Lymphogranuloma venereum

Mycoplasma genitalium

Trichomoniasis

Syndrome-Based Diseases

Progress Tracker

Gonorrhea

14
questionsTreatment of Antimicrobial-Resistant
Gonococcal Infection

1 2 3 4 5 6 7 8 9 10 11 12 13 14

Explore Topic Questions

Learning Objectives

1. Summarize recommendations regarding routine screening for *Neisseria gonorrhoeae* infection in women and men.
2. List CDC-recommended treatment of uncomplicated urogenital and rectal gonococcal infection.
3. State treatment recommendations for persons with pharyngeal gonorrhea.
4. Describe appropriate management of sex partners of persons diagnosed with gonorrhea.
5. Summarize epidemiology and treatment of antimicrobial-resistant *Neisseria gonorrhoeae*.

Authors

Rebecca G. Kinney, MD
Lindley A. Barbee, MD, MPH

CNE/CME Continuing Education

This question bank topic qualifies for **1.5 CME AMA PRA Category 1 Credits™** or **1.5 CNE contact hours**. [View CE Notices](#).**CNE and CME Origination:** February 1st, 2017**CNE and CME Expiration:** January 31st, 2020**Steps to Acquire CE for this Question Bank Topic:**



STD Question Bank

Pathogen-Based
Diseases

Bacterial Vaginosis

Candidiasis – Vulvovaginal

Chancroid

Chlamydia

Gonorrhea

Granuloma Inguinale

Herpes Simplex Virus – Genital

Lymphogranuloma venereum

Mycoplasma genitalium

Trichomoniasis

Syndrome-Based Diseases >

Progress Tracker

Gonorrhea

< 1 2 3 4 5 6 7 8 9 10 11 12 13 14 >

A 21-year-old male patient comes in for a return visit to the STD clinic with urethral discomfort and purulent discharge. Five days prior he was diagnosed with urethral gonorrhea and was treated with recommended therapy. He states his symptoms have not improved and he has not had any type of sexual contact with anyone since receiving treatment for gonorrhea. A diagnosis of antimicrobial-resistant *Neisseria gonorrhoeae* infection is suspected and a urethral swab for culture is obtained.

Which one of the following is the best choice for treating gonorrhea when there is a high suspicion for antimicrobial-resistant *Neisseria gonorrhoeae* infection?

- “High-dose” therapy with ceftriaxone 1 g intramuscular as a single dose plus azithromycin 2 g orally as a single dose
- Moxifloxacin 400 mg orally once daily for 3 days plus doxycycline 100 mg orally twice daily for 7 days
- Gentamicin 240 mg intramuscularly as a single dose plus azithromycin 2 g orally as a single dose
- Clindamycin 600 mg three times daily for 7 days plus atovaquone 1500 mg twice daily for 7 days

< Previous Question

Skip Question >

Check Answer

Progress Not Saved!

Since you are not signed in, your progress won't be saved.

Sign In or Register

Question Last Updated

April 26th, 2017



STD Question Bank

Pathogen-Based
Diseases

Bacterial Vaginosis

Candidiasis – Vulvovaginal

Chancroid

Chlamydia

Gonorrhea

Granuloma Inguinale

Herpes Simplex Virus – Genital

Lymphogranuloma venereum

Mycoplasma genitalium

Trichomoniasis

Syndrome-Based Diseases

Progress Tracker

Gonorrhea

< 1 2 3 4 5 6 7 8 9 10 11 12 13 14 >

A 21-year-old male patient comes in for a return visit to the STD clinic with urethral discomfort and purulent discharge. Five days prior he was diagnosed with urethral gonorrhea and was treated with recommended therapy. He states his symptoms have not improved and he has not had any type of sexual contact with anyone since receiving treatment for gonorrhea. A diagnosis of antimicrobial-resistant *Neisseria gonorrhoeae* infection is suspected and a urethral swab for culture is obtained.

Which one of the following is the best choice for treating gonorrhea when there is a high suspicion for antimicrobial-resistant *Neisseria gonorrhoeae* infection?

You chose this option correctly:

- Gentamicin 240 mg intramuscularly as a single dose plus azithromycin 2 g orally as a single dose

Summary

Evolving drug resistance, particularly to the cephalosporin class, is a significant threat to the treatment of *Neisseria gonorrhoeae*, and clinicians should remain vigilant for cases of treatment failure. If a cephalosporin-resistant isolate is highly-suspected or documented on susceptibility testing, retreatment is recommended with either:

- Azithromycin 2 g orally as a single dose plus gemifloxacin 320 mg orally as a single dose, or
- Azithromycin 2 g orally as a single dose plus gentamicin 240 mg intramuscularly as a single dose

Note that in the United States, there is currently a drug shortage of gemifloxacin related to a legal dispute regarding the license to manufacture and distribute gemifloxacin. Clinicians who are treating patients with suspected or confirmed cephalosporin resistance should also consult an infectious disease expert and report treatment failure to the Centers for Disease Control (CDC) within 24 hours of diagnosis. In cases of retreatment after treatment failure, a test-of-cure is recommended 7 to 14 days after retreatment. In addition, sexual partners of patients who are found to have an antimicrobial-resistant strain of gonorrhea will require treatment with the same regimen as the source patient.

Links to Other National Network of STD Clinical Prevention Training Centers



STD Modules Self-Study >



STD Modules Quick Reference >



Question Bank



Clinical Consultation




Master Bibliography



National STD Curriculum

Contributors

Funded by a grant from the Centers for Disease Control and Prevention



[LINK to: STD Clinical Consultation](#)
Provided by the STD Clinical Consultation Network



STD Clinical Consultation Network


For Licensed Healthcare Professionals and STD Program Staff

- The *National STD Curriculum* does not provide clinical consultation. If you are interested in obtaining clinical consultation regarding STD-related patient care, please contact the STD Clinical Consultation Network.
- The STD Clinical Consultation Network is a free clinical consultation service provided by expert faculty at regional STD Prevention Training Centers, as part of the National Network of STD Clinical Prevention Training Centers.
- The STD clinical consultation service is available only to licensed healthcare professionals and STD program staff.



National Network of
STD Clinical Prevention
Training Centers

STD Clinical Consultation Network

Consultations can be submitted at <https://www.STDCCN.org> 
or by clicking on the logo above.

Benefits for National STD Curriculum members

Signed in members are able to pre-fill some of the required information on the STD CCN consultation form.

Already have an account? [Sign in](#) to take advantage of this benefit.

www.STDCCN.org

About the National STD Curriculum

The *National STD Curriculum* is a free educational Web site from the University of Washington STD Prevention Training Center and the University of Washington. This project is funded by a grant from the Centers for Disease Control and Prevention (CDC).

The *National STD Curriculum* addresses the epidemiology, pathogenesis, clinical manifestations, diagnosis, management, and prevention of STDs.

Free [CME](#) credit and free [CNE](#) credit are offered throughout the site.

Contributors

Site Overview

News

STD Question Bank with CE

The STD Question Bank features interactive board-review style questions that emphasize key points in the 2015 STD Treatment Guidelines. [CNE/CME](#)

Free CE Available! [CNE/CME](#)

Currently available via the Self-Study Modules. Click for details.



About the National Network of STD Clinical Prevention Training Centers

The *National STD Curriculum* is part of the National Network of STD Clinical Prevention Training Centers (NNPTC), a CDC-funded group of regional and national STD training centers created in partnership with health departments and universities.

The NNPTC is dedicated to increasing the knowledge and skills of health professionals by providing a spectrum of state-of-the-art trainings for STD prevention and treatment with a particular focus on experiential learning.

To learn more about the NNPTC and STD trainings in your region, please visit the NNPTC web site at <http://nnptc.org/>.

Funded by a grant from the Centers for Disease Control and Prevention

UNIVERSITY of WASHINGTON

THE UNIVERSITY OF ALABAMA AT BIRMINGHAM

Self-Study Modules [CNE/CME](#)

Chlamydia
Gonorrhea
Herpes Simplex Virus (HSV)
Human Papillomavirus (HPV)
Pelvic Inflammatory Disease (PID)
Syphilis
Vaginitis

Quick Reference

Chlamydia
Gonorrhea
Herpes Simplex Virus (HSV)
Human Papillomavirus (HPV)
Pelvic Inflammatory Disease (PID)
Syphilis
Vaginitis

Question Bank [CNE/CME](#)

Case Series
Clinical Consultation
Master Bibliography

Contributors
Site Overview
Content Bundles





National Network of STD Clinical Prevention Training Centers



ABOUT

TRAINING CENTERS

TRAINING

STD CONSULTATION

NEWS

RESOURCES


FTCC

ARE YOU READY TO **DISCUSS,**
DIAGNOSE, TREAT, MANAGE,
REFER & PREVENT STDS?



www.nnptc.org

Synergy of Three CDC-Funded STD Programs



Master Bibliography

Modules

Pathogen-Based Diseases **1**

Syndrome-Based Diseases **2**

TIP! Use the Shift key or the checkbox to select more than one module at a time

Topics:

1.1. Bacterial Vaginosis

1.1.A. Additional References

1.2. Candidiasis – Vulvovaginal

1.2.A. Additional References

1.3. Chancroid

1.3.A. Additional References

1.4. Chlamydia

1.4.1. Epidemiology in the United States

1.4.2. Microbiology and Pathogenesis

1.4.3. Clinical Manifestations

1.4.4. Laboratory Diagnosis

1.4.5. Screening for Chlamydial Infection

1.4.6. Treatment

1.4.7. Patient Counseling and Education

1.6. Gonorrhea

1.6.1. Epidemiology in the United States

1.6.2. Microbiology and Pathogenesis

1.6.3. Clinical Manifestations

1.6.4. Laboratory Diagnosis

1.6.5. Screening for Gonococcal Infection

1.6.6. Treatment

1.6.A. Additional References

View by Topic

View by Author

Search

Bibliography

The bibliography lists all references used in the online course. The View by Topic selection organizes the references sequentially by Module, Lesson, and subsections in the lessons. The View by Author lists the authors in alphabetical order.

Disease Type 1. Pathogen-Based Diseases

Disease 1. Bacterial Vaginosis

Additional References

Abad CL, Safdar N. The role of lactobacillus probiotics in the treatment or prevention of urogenital infections--a systematic review. *J Chemother.* 2009;21:243-52.

[PubMed Abstract](#)

Allsworth JE, Peipert JF. Prevalence of bacterial vaginosis: 2001-2004 National Health and Nutrition Examination Survey data. *Obstet Gynecol.* 2007;109:114-20.

[PubMed Abstract](#)

Amsel R, Totten PA, Spiegel CA, Chen KC, Eschenbach D, Holmes KK. Nonspecific vaginitis. Diagnostic criteria and microbial and epidemiologic associations. *Am J Med.* 1983;74:14-22.

[PubMed Abstract](#)

Antonio MA, Meyn LA, Murray PJ, Busse B, Hillier SL. Vaginal colonization by probiotic *Lactobacillus crispatus* CTV-05 is decreased by sexual activity and endogenous Lactobacilli. *J Infect Dis.* 2009;199:1506-13.

[PubMed Abstract](#)

Balkus JE, Manhart LE, Lee J, et al. Periodic Presumptive Treatment for Vaginal Infections May Reduce the Incidence of Sexually Transmitted Bacterial Infections. *J Infect Dis.* 2016;213:1932-7.

[PubMed Abstract](#)

Bautista CT, Wurapa E, Sateren WB, Morris S, Hollingsworth B, Sanchez JL. Bacterial vaginosis: a synthesis of the literature on etiology, prevalence, risk factors, and relationship with chlamydia and gonorrhea infections. *Mil Med Res.* 2016;3:4.

[PubMed Abstract](#)

Beigi RH, Austin MN, Meyn LA, Krohn MA, Hillier SL. Antimicrobial resistance associated with the treatment of bacterial vaginosis. *Am J Obstet Gynecol.* 2004;191:1124-9.

[PubMed Abstract](#)

Bilardi JE, Walker S, Temple-Smith M, et al. The burden of bacterial vaginosis: women's experience of the physical, emotional, sexual and social impact of living with recurrent bacterial vaginosis. *PLoS One.* 2013;8:e74378.

[PubMed Abstract](#)

More than 1200 STD-related references

Accounts, Tracking Progress, and CE



STD Modules Self-Study >


STD Modules Quick Reference >

Question Bank

Clinical Consultation

Master Bibliography


Self-Study STD Modules




Chlamydia



Gonorrhea



HSV




HPV



PID



Syphilis



Vaginitis

Self-Study STD Modules for Clinicians

You are just a few steps away from free CE credits!

1 ★

Sign in or Register

A free account is required.

2

Study the Materials

An entire module, or just a few topics at a time.

3

Take the CE Quiz

5 questions covering the topics in each module.

4

Claim CE Credit

Free CNE and Free CME available!

New Users

Create a free account to get started.

Required for CE

Register >

Returning Users

Email Address

Password

[Forgot password?](#)

Sign in >



**Account Management****Register**

as an Individual »

as an Organization »

Account Retrieval

Join a Group

Register

Why create an account?

- Creating a FREE account will give you access to the Self-Study Modules
- You will be able to claim CE credits **CME/CNE**
- You will have the ability to resume this curriculum at any point in time

➔ **The fields marked by a red arrow are required.**

Fields marked by a ✨ are displayed on your Certificate of Completion.

➔ **Account Type**

Please select the account type that best matches you:

Individual Account

You are an Individual and would like to:

- Track progress for yourself or collect CE credits
- Resume the curriculum at a later time from any internet connected computer
- Join a group or an organization (only by invitation)

Organization/Group Account

You are a Teacher, Coordinator, Manager or Director and would like to:

- Track progress of your staff or students



Welcome Back!

You were last online on January 20th, 2018 at 11:51 am.



Self-Study STD Modules



Chlamydia ✓ Gonorrhea ✓ HSV ✓ HPV ✓ PID ✓ Syphilis ✓ Vaginitis ✓ Certificate of Completion

Progress Tracker - 1st Edition

CNE Certificate »

	Chlamydia	Gonorrhea	HSV	HPV	PID	Syphilis	Vaginitis
Topic 1	✓	✓	✓	✓	✓	✓	✓
Topic 2	✓	✓		✓	✓	✓	✓
Topic 3	✓	✓		✓	✓	✓	✓
Topic 4	✓	✓		✓	✓	✓	✓
Topic 5	✓	✓	✓		✓	✓	✓
Topic 6	✓	✓	✓		✓	✓	✓
Topic 7	✓	✓			✓	✓	
Topic 8	✓	✓	✓			✓	
Topic 9						✓	
Topic 10							
CE Quiz	✓ CNE ✓	✓ CNE ✓	✓ CNE ✓	✓	✓	✓	✓





Self-Study STD Modules

- Home
- Chlamydia
- Gonorrhea
- HSV
- HPV
- PID
- Syphilis
- Vaginitis
- Certificate of Completion

🌟 Your Certificate of Completion

CE (CNE and CME) Certificates CNE/CME

You can earn CNE or CME for each module in this set. For CNE, your certificate will be provided by the University of Washington School of Nursing. For CME, your certificate will be provided by the University of Alabama at Birmingham. Each of these certificates will reflect the number of modules completed.

CE As You Go! You earn CE as you complete each module, and DO NOT have to complete all modules before printing your CE certificate.

Print your CE certificate from the Home screen or one of the modules where you obtained that CE.

Non-CE Certificate

This certificate only shows the self-study modules you complete and does not show continuing education credits.



Print Certificate

Your certificate will show your progress thus far.

View Certificate



Complete Modules

Each module completed will be displayed on your certificate.

7/7
modules completed



STD Question Bank

Hello [User Name]

The STD Question Bank features interactive board-review style questions that emphasize key points in the 2015 STD Treatment Guidelines.

Your progress and CE credits earned in the question bank will be reflected in the following certificates:

Free CNE/CME Available!
Pick a topic below to begin.

CNE Certificate

Score 80% or higher in a topic for the Non-CE certificate

Non-CE Certificate of Completion

MY PROGRESS

MY DASHBOARD

MY GROUPS

Tip The Progress Tracker gives you an overview of all of the questions. You can view it on your dashboard, or access it from any other page from the bottom of the screen. You can also access it from the User menu at the top of the page.

Tip Use the Folder icon next to a topic to compare sets of answers. The icon only shows up for topics which have multiple sets of answers.

CE A tag will be displayed next to topics for which CE is claimed. **CNE/CME** displayed next to the score means that's the set of answers during which the CE was earned.

Pathogen-Based Diseases

Bacterial Vaginosis	1 2 3 4 5 6 7	
Candidiasis - Vulvovaginal	1 2 3 4 5 6 7 8	13%
Chancroid	1 2 3 4 5	CE Earned 100%
Chlamydia	1 2 3 4 5 6 7 8 9 10 11	45%
Gonorrhea	1 2 3 4 5 6 7 8 9 10 11 12 13 14	
Granuloma Inguinale	1 2 3 4 5	60%
Herpes Simplex Virus - Genital	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17	0%
Lymphogranuloma venereum	1 2 3 4 5 6	CE Earned 83%
<i>Mycoplasma genitalium</i>	1 2 3 4 5	CE Earned 80%
Trichomoniasis	1 2 3 4 5 6 7 8	13%

Tip The Progress Tracker gives you an overview of all of the questions. You can view it on your dashboard, or access it from any other page from the bottom of the screen. You can also access it from the User menu at the top of the page.

Tip Use the Folder icon next to a topic to compare sets of answers. The icon only shows up for topics which have multiple sets of answers.

CE A tag will be displayed next to topics for which CE is claimed. **CNE/CME** displayed next to the score means that's the set of answers during which the CE was earned.


Pathogen-Based Diseases

Bacterial Vaginosis	1 2 3 4 5 6 7	
Candidiasis - Vulvovaginal	1 2 3 4 5 6 7 8	13%
Chancroid	1 2 3 4 5	CE Earned 100%
Mar 27, 2017, 10:10 am	1 2 3 4 5	60%
Mar 27, 2017, 9:46 am	1 2 3 4 5	20%
Chlamydia	1 2 3 4 5 6 7 8 9 10 11	45%
Gonorrhea	1 2 3 4 5 6 7 8 9 10 11 12 13 14	
Granuloma Inguinale	1 2 3 4 5	60%
Herpes Simplex Virus - Genital	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16 17	0%
Lymphogranuloma venereum	1 2 3 4 5 6	CE Earned 83%
<i>Mycoplasma genitalium</i>	1 2 3 4 5	CE Earned 80%
Trichomoniasis	1 2 3 4 5 6 7 8	13%
Syndrome-Based Diseases		
Epididymitis	1 2 3 4 5 6 7 8 9 10 11	
Proctitis, Proctocolitis, and Enteritis	1 2 3 4 5 6	0%



Certificate of Completion

University of Washington and the National STD Curriculum
certify that



has completed the

Chlamydia, Gonorrhea, HSV, HPV, PID, Syphilis, and Vaginitis

modules of the

Self-Study STD Modules



David H. Spach, MD
Professor of Medicine
University of Washington
Editor-in-Chief
National STD Curriculum

Certificate Details

This certificate shows the progress of **Andrew Karpenko, BS**, a Software Engineer, identified by akarpy@uw.edu as of January 20, 2018, 12:05 pm.
You may validate this certificate or obtain a digital copy by visiting <https://depts.washington.edu/ptostd/dev/certificate/self-study/akarpy@uw.edu>

The following Modules were completed:

Modules	Learning Objectives	Completed as of
Chlamydia	<ol style="list-style-type: none">1. Summarize the epidemiology of <i>Chlamydia trachomatis</i> infections in the United States2. Describe the microbiology, life cycle, and transmission of <i>Chlamydia trachomatis</i>.3. Discuss the clinical manifestations of chlamydial infections in men, women, and children.4. Compare the common laboratory diagnostic methods used to diagnose chlamydial infections.5. State routine chlamydial screening recommendations for different patient populations.6. List the CDC-recommended treatment regimens for chlamydial infections.7. Summarize counseling and education messages for individuals with chlamydial infection.	May 22, 2017



Self-Study STD Modules



Chlamydia

Gonorrhea

HSV

HPV

PID

Syphilis

Vaginitis



Certificate of Completion

Module 1. Chlamydia - 100% complete

Steps to Acquire CE for this Activity:

1 ✓

CE Quiz

🟢 CNE earned

2 ✓

Provide Feedback

Survey Completed

3 ★

Print Certificate



Certifies that



has earned CME credit for the following activities:

National STD Curriculum

Question Bank

12/05/2017	Granuloma Inguinale	0.5
	<i>Total AMA PRA Category 1 Credits™</i>	0.5

STD Self-Study Modules

011/26/2017	Herpes Simplex Virus – Genital	1
011/16/2017	Human Papillomavirus Infection	1
	<i>Total AMA PRA Category 1 Credits™</i>	2

Funded by the Centers for Disease Control and Prevention, the University of Washington developed the National STD Curriculum, the Hepatitis C Online, and the Hepatitis B Web Study websites to enhance American healthcare professionals' ability to diagnose, treat and manage these diseases. For the same purpose, the U.S. Health Resources & Services Administration funded the University of Washington to develop the National HIV Curriculum.


For information on applicability and acceptance of continuing education credit(s) for your work, please consult your professional licensing board.

Ronan O'Beirne, EdD
 Director, Continuing Medical Education

1600 Hill, Suite 1100
 1570 University Blvd
 Birmingham, AL 35293
 205.934.2607
 www.uab.edu/med

The University of
 Alabama at Birmingham
 1615 6th Ave. S.
 Birmingham, AL 35294-0001

National STD Curriculum Learning Groups



Director/Administrator/Leader

**Account Management****Register**

as an Individual »

as an Organization »

Account Retrieval

Join a Group

Register

Why create an account?

- Creating a FREE account will give you access to the Self-Study Modules
- You will be able to claim CE credits **CME/CNE**
- You will have the ability to resume this curriculum at any point in time

➔ **The fields marked by a red arrow are required.**

Fields marked by a ✨ are displayed on your Certificate of Completion.

➔ **Account Type**

Please select the account type that best matches you:

Individual Account

You are an Individual and would like to:

- Track progress for yourself or collect CE credits
- Resume the curriculum at a later time from any internet connected computer
- Join a group or an organization (only by invitation)

Organization/Group Account

You are a Teacher, Coordinator, Manager or Director and would like to:

- Track progress of your staff or students

UW Testing Group

HOME

PROGRESS

ANALYTICS

MEMBERS

SETTINGS

Only current members' progress is included. Use the tabs below to look at progress within the individual parts of the curriculum.

Active Members (last 30 days)

Current Members

COURSE MODULES

QUESTION BANK

View results for:

Self-Study STD Modules

Chlamydia Gonorrhea Herpes Simplex Virus (HSV) Human Papillomavirus (HPV) Pelvic Inflammatory Disease (PID) Syphilis Vaginitis

	Chlamydia	Gonorrhea	Herpes Simplex Virus (HSV)	Human Papillomavirus (HPV)	Pelvic Inflammatory Disease (PID)	Syphilis	Vaginitis	
1								Progress
								Score
2	53 100	100 ✓	40	0	25	80	60 100	Progress Score
3	100 ✓ 100	100 ✓	100	100 ✓	100 ✓	100 ✓	100 ✓	Progress Score
4				20				Progress Score
5	100 ✓ 100	62 100	20					Progress Score
6								Progress Score
7								Progress Score

Key

- CNE/CME - CE was earned
 - CNE/CME - CE was attempted
- User progress is delayed by up to 15 minutes

UW Testing Group

- HOME
- PROGRESS**
- ANALYTICS
- MEMBERS
- SETTINGS

Only current members' progress is included. Use the tabs below to look at progress within the individual parts of the curriculum.

Active Members (last 30 days) Current Members

COURSE MODULES

QUESTION BANK

Pathogen-Based Diseases Syndrome-Based Diseases

	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	1	2	3	4	5	6	7	
1	100			100																			Progress
	86			100																			Score
2	100		100																				Progress
	86		100																				Score
3		100		100			100					100	100										Progress
		100		100			100					83	80										Score
4																							Progress
																							Score
5		50																					Progress
		50																					Score
6	100	100	100	100		93	100		100			100	100		89		100				100		Progress
	100	100	60	91		93	80		71			83	60		100		82				100		Score
7																							Progress
																							Score

Key


- CNE/CME - CE was earned
 - CNE/CME - CE was attempted
- User progress is delayed by up to 15 minutes

NATIONAL STD CURRICULUM

“1 Year” Data

National STD Curriculum

Number of Registered Users by Month




National STD Curriculum

Cumulative Number of Registered Users




National STD Curriculum

Running Sum of Modules + Question Bank Completed




National STD Curriculum Running Sum Continuing Education Credits (CNE + CME)



Check on Learning Activity in Self Study Modules


Check on Learning Activity in Self-Study Modules			
Title of Module	# Questions	# Participants	# Answers
Chlamydia	12	6,986	72,046
Gonorrhea	10	5,182	46,814
Herpes Simplex Virus	9	4,426	35,995
Human Papillomavirus	10	3,873	34,856
PID	10	3,292	29,973
Syphilis	10	3,578	31,072
Vaginitis	13	3,726	43,035
TOTAL	74		293,791

National STD Curriculum Registered Users by Zip Code




National STD Curriculum Registered Users by Zip Code


STD Curriculum Registered Users




CT Rate by county



GC Rate by county



Syphilis Rate by county



NATIONAL STD CURRICULUM

Future Developments for Curriculum

National STD Curriculum Expansion of Question Bank

STD Question Bank

The STD Question Bank features interactive board-review style questions that emphasize key points in the 2015 STD Treatment Guidelines.

Free CNE/CME Available!
Pick a topic below to begin.

New Users
Create a free account to get started.
Required for CE
Register

Returning Users National STD Curriculum
Email Address
Password
Forgot password? Sign In

Explore the Questions
All of the questions are available for free. Start learning and reviewing material anytime without the need to register.
Choose a topic from the list to begin

Track Progress CNE/CME
Save your progress! Creating an account and signing in allows you to track questions you have answered, see your progress and resume at a later date. An account is required for CE.
Sign In or Register to Track Progress

CDC Centers for Disease Control and Prevention
CDC 24/7: Saving Lives, Protecting People™

2015 Sexually Transmitted Diseases Treatment Guidelines

CONTENTS

Introduction	1
Methods	1
Clinical Prevention Guidance	2
Special Populations	9
Emerging Issues	17
Hepatitis C	17
<i>Mycoplasma genitalium</i>	20
HIV Infection: Detection, Counseling, and Referral	21
Diseases Characterized by Genital, Anal, or Perianal Ulcers	25
Chancroid	26
Genital HSV Infections	27
Granuloma Inguinale (Donovanosis)	32
Lymphogranuloma Venereum	33
Syphilis	34
Management of Persons Who Have a History of Penicillin Allergy	49
Diseases Characterized by Urethritis and Cervicitis	51
Urethritis	51
Nongonococcal Urethritis	52
Cervicitis	53
Chlamydial Infections	55

CONTENTS (Continued)

Gonococcal Infections	60
Diseases Characterized by Vaginal Discharge	69
Bacterial Vaginosis	69
Trichomoniasis	72
Vulvovaginal Candidiasis	75
Pelvic Inflammatory Disease	78
Epididymitis	82
Human Papillomavirus Infection	84
Anogenital Warts	86
HPV-Associated Cancers and Precancers	90
Viral Hepatitis	94
Hepatitis A	94
Hepatitis B	95
Proctitis, Proctocolitis, and Enteritis	100
Ectoparasitic Infections	101
Pediculosis Pubis	101
Scabies	102
Sexual Assault and Abuse and STDs	104
References	110
Terms and Abbreviations Used in This Report	135

STD Treatment Guidelines Learning and Assessment Tool

National STD Curriculum: STD Case Studies Revision

Tentative Planned Topics


- Routine STD history and examination
- Genital ulcer
- Cervicitis
- Vaginal discharge
- Urethritis/Urethral discharge
- Proctitis
- Testicular swelling and pain
- Evaluation and management of acute HIV

National STD Curriculum


Clinical Challenges/Expert Opinions

- Expert opinions for challenging cases
- Expert opinions for controversial topics

National STD Curriculum Power Point Slide Decks




Primary and Secondary Syphilis — Rates of Reported Cases by Age and Sex, U.S., 2016




Source: CDC. Sexually Transmitted Disease Surveillance 2016. Syphilis.

Syphilis — Reported Cases of Congenital Infection, United States, 1941–2016






Source: CDC. Sexually Transmitted Disease Surveillance 2016. Syphilis.

Tube Dilution with Syphilis RRP Testing



Sample Syphilis VDRL Titer Changes



National STD Curriculum

Long-Range Plans: Learning Platform/Portfolio

STD Modules
Self Study

Question Bank

Case Studies

Clinical
Challenges

Power Point
Slide Decks



STD Modules Self-Study >



STD Modules Quick Reference >



Question Bank



Clinical Consultation




Master Bibliography



National STD Curriculum

Contributors

Funded by a grant from the Centers for Disease Control and Prevention



Visit the *National STD Curriculum* at:
www.std.uw.edu

Questions?
Use chat box